

Localization of Gravitino Field on Thin Branes

Yun-Zhi Du^{1a}, Li Zhao^{1b}, Xiang-Nan Zhou^{2c}, Yi Zhong^{1d}, and Yu-Xiao Liu^{1e}

¹*Institute of Theoretical Physics, Lanzhou University, Lanzhou 730000, People's Republic of China*

²*College of Physics and Information Engineering,
Shanxi Normal University, Linfen 041004, People's Republic of China*

In this paper, we investigate the localization of a gravitino field on two kinds of thin branes, the Randall-Sundrum-1 (RS1) branes and the scalar-tensor branes. The coupled chiral equations for the Kaluza-Klein (KK) modes of a gravitino field are obtained by the gravitino chiral KK decompositions with the supersymmetry transformations. It is shown that, in the RS1 model for the left- and right-handed zero modes of the gravitino field, only one of them can be localized near one brane. For the massive modes, both chiral modes survive and the lower KK modes are localized near the IR brane from the four-dimensional physical coordinate point of view. For the scalar-tensor brane model, the localization of the gravitino chiral zero modes depends on the coupling parameter λ , and they will be not localized around anyone brane within a certain range of the parameter λ , which is quite different from the RS1 model. Furthermore, we also give the corresponding mass spectra of the massive KK gravitinos in the scalar-tensor model.

arXiv:1511.03102v1 [hep-th] 10 Nov 2015

^a duyzh13@lzu.edu.cn

^b lizhao@lzu.edu.cn, corresponding author

^c zhouxn10@lzu.edu.cn

^d zhongyi11@lzu.edu.cn

^e liuyx@lzu.edu.cn, corresponding author

I. INTRODUCTION

It is well-known that our visible world consists of one-dimensional time and three-dimensional space. However, the Kaluza-Klein (KK) theory [1, 2] and string theory indicate that there are extra spacial dimensions beyond our familiar ones. Inspired by string theory, Antoniadis, Arkani-Hamed, Dimopoulos, and Dvali suggested that our four-dimensional universe is a membrane embedded in a higher-dimensional spacetime with flat large extra dimensions [3, 4]. Subsequently, Randall and Sundrum proposed a warped braneworld scenario in a five-dimensional Anti-de Sitter (AdS) spacetime, where the extra dimension is warped [5, 6]. The braneworld theory has attracted much attention because it can provide an alternative mechanism to solve the hierarchy problem (the 16 order difference between the weak scale and the Planck scale). Furthermore, extra dimensions in such theory may be probed by current experiments [7–15]. This is different from the original KK theory and string theory, where extra dimensions are very hard to be detected by current or near future experiments. Some of the possible experimental probes of extra spatial dimensions include the possible creation of mini-black holes [16–18] and the collider cross sections for interactions between the Standard Model particles and KK particles [19–21] at the Large Hadron Collider as well as deviations of Newton’s law of gravity at the micro-meter or smaller distance in current table top experimental searches [22–26].

In the braneworld scenario, all particles/fields in the Standard Model should be localized on a $(3 + 1)$ –dimensional brane. Therefore, we need localization mechanisms for different spin fields to make our world appears effectively $(3 + 1)$ –dimensional up to some energy scale. Localization of various matters in some kinds of braneworld models has been extensively investigated for examples in Refs. [27–47]. In this paper, we are interested in the localization of a five-dimensional massive gravitino field on thin branes.

Gravitino is the particle with spin $3/2$. It was proposed as the supersymmetric partner of graviton in the supergravity theory. In particular, this particle is regarded as a candidate of dark matter in cosmology [48–53]. It was shown that if the axino mass is around 1 MeV and gravitino mass around 1 eV, then hot axions and gravitinos could act like hot dark matters [48]. In Ref. [50], gravitino is a promising candidate for the cold dark matter with new bounds on its mass. However, the light gravitinos with mass in 1 eV to 1 MeV range as the dark matter candidate are muddled by early-universe uncertainties [54]. The authors in Ref. [54] had explained how upcoming data from colliders may clarify this uncertainty picture. Furthermore, in the supersymmetric Standard Models the lightest supersymmetric particle is an ultralight gravitino ($m_{3/2} \sim 1$ eV). A simple test for the lightness of gravitino at the Large Hadron Collider was proposed in Ref. [55]. Recently, some attention was paid to gravitino fields in black hole backgrounds [56–59]. Therefore, the localization and mass spectrum of a gravitino field on a $(3 + 1)$ –dimensional brane are of great significance in the brane scenario.

The localization of a gravitino field (Ψ_M) was investigated in Refs. [29, 31, 35, 60–62]. Just like the case of a free Dirac fermion, a free gravitino cannot be confined to a RS-like brane in a five-dimensional spacetime. Thus, the authors of Refs. [31, 60] studied the localization of zero mode of a five-dimensional bulk gravitino field by introducing a bulk mass term. It was shown in Refs. [35, 61] that, by taking some gauges such as $\Psi_5 = 0$, the zero mode of an interacting bulk gravitino in a D –dimensional spacetime ($D \geq 6$) has similar localization property as that of an interacting bulk Dirac fermion. In a six-dimensional gauged supergravity model, the brane-induced gravitino mass term would lead to a power-like suppressed on the effective four-dimensional gravitino mass [62]. Furthermore, the coupling of the gravitino KK modes with other fermions and with their scalar supersymmetrical partners on the brane was investigated in Ref. [63]. In this paper, we would like to consider a five-dimensional massive gravitino field in the RS1 and scalar-tensor brane models [43] through the supersymmetric transformations of this field, instead of taking the gauge $\Psi_5 = 0$. Our aim is to study the localization of chiral zero modes and the mass spectrum of KK gravitinos.

This work is organized as follows. In Sec. II, we consider a massive gravitino field in a five-dimensional spacetime with the extra dimension compactified on S^1/Z_2 and introduce the supersymmetric transformations of this field. In particular, the coupled chiral equations and the orthonormality conditions of the left- and right-handed KK gravitinos are derived. The localization and mass spectrum of the gravitino field on the RS1 branes and the scalar-tensor branes are investigated in Sec. IV and Sec. III, respectively. Finally, the discussion and conclusion are presented in Sec. V.

II. LOCALIZATION OF A GRAVITINO FIELD ON THIN BRANES

In this section, we consider the localization of a gravitino field on a thin brane embedded in a five-dimensional spacetime with coordinates $x^M = (x^\mu, y)$. Here, the five-dimensional spacetime indices are denoted by $M, N, \dots = 0, 1, 2, 3, 5$ and the four-dimensional Minkowskian spacetime indices $\mu, \nu, \dots = 0, 1, 2, 3$. To avoid deviation from the detection by current experiments, one can think that the extra dimension y is finite and the simplest way is to assume a periodic geometry of the extra dimension such as a circle S^1 with radius R . However, the chiral fermions in the Standard Model cannot be described by compactifying bulk fermions on a circle. Instead the fifth dimension can be compactified on a line segment S^1/Z_2 with Z_2 representing the identification $y \leftrightarrow -y$ (so we have $0 \leq y \leq \bar{y} = \pi R$).

This is the well-known Z_2 orbifold symmetry [6, 43]. There are two thin 3-branes located at two orbifold fixed points $y = 0$ and $y = \bar{y}$.

The line element of the five-dimensional spacetime is given by

$$ds^2 = e^{-2\sigma} \eta_{\mu\nu} dx^\mu dx^\nu + dy^2, \quad (1)$$

where the warped factor $e^{-2\sigma}$ is only the function of $|y|$. Based on the metric (1), the warped five-dimensional vielbeins $e_M^{\bar{M}}$ and the flat four-dimensional ones $\hat{e}_\mu^{\bar{\mu}}$ read

$$e_M^{\bar{M}} = \begin{bmatrix} e^{-\sigma} & \hat{e}_\mu^{\bar{\mu}} & 0 \\ 0 & & 1 \end{bmatrix}, \quad e_M^{\bar{M}} = \begin{bmatrix} e^\sigma & \hat{e}^{\mu\bar{\mu}} & 0 \\ 0 & & 1 \end{bmatrix}, \quad \hat{e}_\mu^{\bar{\mu}} = \begin{bmatrix} 1 & & \\ & 1 & \\ & & 1 \end{bmatrix}. \quad (2)$$

Here, $\bar{M}$ and $\bar{\mu}$ denote the (4+1)-dimensional and (3+1)-dimensional local Lorentz indices, respectively. With the relations $e_{M\bar{M}} = g_{MN} e_N^{\bar{M}}$ and $e^{M\bar{M}} = g^{MN} e_N^{\bar{M}}$, we have

$$e_{M\bar{M}} = \begin{bmatrix} e^{-\sigma} & \hat{e}_{\mu\bar{\mu}} & 0 \\ 0 & & 1 \end{bmatrix}, \quad e^{M\bar{M}} = \begin{bmatrix} e^\sigma & \hat{e}^{\mu\bar{\mu}} & 0 \\ 0 & & 1 \end{bmatrix}, \quad \hat{e}_{\mu\bar{\mu}} = \eta_{\bar{\mu}\bar{\nu}} \hat{e}_\mu^{\bar{\nu}}. \quad (3)$$

In Refs. [64–68], the authors gave the complete supergravity action in the RS1 model, which includes the gravitino and graviphoton together with the graviton. In an AdS_5 spacetime, the graviphoton is set to zero [64]. In order to study the supersymmetry breaking, the additional gravitino kinetic and mass terms were considered in Ref. [60].

It is known that the Dirac fermion mass parameter is odd under the Z_2 transformation in a five-dimensional spacetime with the extra spacial dimension compactified on S^1/Z_1 , so is the gravitino mass parameter. Therefore, for the case of a gravitino field, the simplest additional mass term can be proportional to $\sigma' \equiv \partial_y \sigma$. Furthermore, the gravitino mass arises from an odd ‘kink’ profile background scalar field with a vacuum expectation value and the vacuum configuration of the scalar field can be regarded as an infinitely thin domain wall. Hence, we adopt the action for a gravitino field with an additional mass term in the five-dimensional spacetime as [60]

$$S_{\frac{3}{2}} = \int d^5x \sqrt{-g} \left(\bar{\Psi}_M i \Gamma^{[MNL]} D_N \Psi_L - a \sigma' \bar{\Psi}_M i \Gamma^{[MN]} \Psi_N \right), \quad (4)$$

where the parameter a is a dimensionless coefficient and $\Gamma^{[MNL]} = \Gamma^M \Gamma^N \Gamma^L$. The matrices Γ^M provide a four-dimensional representation of the Dirac matrices (γ^μ and γ^5) with the chiral representation in a five-dimensional warped spacetime. The covariant derivation $D_N \Psi_L$ is defined as

$$D_N \Psi_L \equiv \partial_N \Psi_L - \Gamma_{NL}^M \Psi_M + \omega_N \Psi_L \quad (5)$$

with the spin connection $\omega_N = \frac{1}{4} \omega_N^{\bar{N}\bar{L}} \Gamma_{\bar{N}} \Gamma_{\bar{L}}$. From the formulas (2) and (3), the relation between the warped gamma matrices Γ^M and the Minkowskian ones $\Gamma^{\bar{M}} = (\Gamma^{\bar{\mu}}, \Gamma^{\bar{5}}) = (\gamma^{\bar{\mu}}, \gamma^{\bar{5}})$ is given by $\Gamma^M = e_M^{\bar{M}} \Gamma^{\bar{M}}$, i.e.,

$$\Gamma^\mu = e^\sigma \gamma^\mu, \quad \Gamma^{\bar{5}} = \gamma^{\bar{5}}. \quad (6)$$

The commutation relations of two gamma matrices are $[\Gamma^M, \Gamma^N] = 2g^{MN}$ and $[\Gamma^{\bar{M}}, \Gamma^{\bar{N}}] = 2\eta^{\bar{M}\bar{N}}$, respectively.

With the metric (1), we have $\omega_\mu = -\frac{1}{2} \sigma' e^{-\sigma} \gamma_\mu \gamma_5$ and $\omega_5 = 0$. Then the nonvanishing components of the covariant derivative $D_N \Psi_L$ can be calculated as follows:

$$D_\mu \Psi_\nu = \partial_\mu \Psi_\nu - \sigma' e^{-2\sigma} \eta_{\mu\nu} \Psi_5 - \frac{\sigma'}{2} e^{-\sigma} \gamma_\mu \gamma_5 \Psi_\nu, \quad (7)$$

$$D_\mu \Psi_5 = \partial_\mu \Psi_5 + \sigma' \Psi_\mu - \frac{\sigma'}{2} e^{-\sigma} \gamma_\mu \gamma_5 \Psi_5, \quad (8)$$

$$D_5 \Psi_\mu = \partial_y \Psi_\mu + \sigma' \Psi_\mu, \quad (9)$$

$$D_5 \Psi_5 = \partial_y \Psi_5 + \sigma' e^{-2\sigma} \Psi_5. \quad (10)$$

Next, we try to give the supersymmetry transformation of the gravitino field described by the action (4), instead of directly taking the gauge $\Psi_5 = 0$ as did in Ref. [31]. Because of the arbitrary parameter a in the action (4), we assume the gravitino supersymmetry transformation as

$$\delta \Psi_M = D_M \eta + b \sigma' \Gamma_M \eta, \quad (11)$$

where b is a parameter, η is the five-dimensional supersymmetry spinor field, and $D_M \eta = \partial_M \eta + \omega_M \eta$. The explicit form of the above supersymmetry transformation for the background metric (1) reads

$$\delta \Psi_\mu = \partial_\mu \eta - \frac{\sigma'}{2} e^{-\sigma} \gamma_\mu \gamma_5 \eta + b \sigma' e^{-\sigma} \gamma_\mu \eta, \quad (12)$$

$$\delta \Psi_5 = \partial_y \eta + b \sigma' \gamma_5 \eta. \quad (13)$$

We can decompose these five-dimensional fields Ψ_M and η into four-dimensional Kaluza-Klein modes [60]

$$\Psi_\mu(x, y) = \sum_n \left[\psi_{L\mu}^{(n)}(x) \xi_{Ln}(y) + \psi_{R\mu}^{(n)}(x) \xi_{Rn}(y) \right], \quad (14)$$

$$\Psi_5(x, y) = \sum_n \left[\psi_{L5}^{(n)}(x) f_{Ln}(y) + \psi_{R5}^{(n)}(x) f_{Rn}(y) \right], \quad (15)$$

$$\eta(x, y) = \sum_n \left[\eta_L^{(n)}(x) \xi_{Ln}(y) + \eta_R^{(n)}(x) \xi_{Rn}(y) \right]. \quad (16)$$

For $M = \mu$, plugging above decompositions into the supersymmetry transformation (12), we have

$$\delta \psi_{L\mu}^{(n)} = \partial_\mu \eta_L^{(n)} + \gamma_\mu \sum_m A_{mn} \eta_R^{(m)}, \quad A_{mn} \equiv \int dy \left(b + \frac{1}{2} \right) \sigma' e^{-2\sigma} \xi_{Rm} \xi_{Ln}, \quad (17)$$

$$\delta \psi_{R\mu}^{(n)} = \partial_\mu \eta_R^{(n)} + \gamma_\mu \sum_m B_{mn} \eta_L^{(m)}, \quad B_{mn} \equiv \int dy \left(b - \frac{1}{2} \right) \sigma' e^{-2\sigma} \xi_{Lm} \xi_{Rn}. \quad (18)$$

The coefficient A_{mn} (B_{mn}) indicates that the supersymmetry transformation of $\psi_{L\mu}^{(n)}$ ($\psi_{R\mu}^{(n)}$) at the level n depends on the whole tower of KK modes $\eta_R^{(m)}$ ($\eta_L^{(m)}$). As $M = 5$, the supersymmetry transformation (13) becomes

$$\delta \Psi_{L5} = \sum_n \delta \psi_{L5}^{(n)} f_{Ln} = \sum_n \left(\eta_L^{(n)} \partial_y \xi_{Ln} + b \sigma' \eta_L^{(n)} \xi_{Ln} \right), \quad (19)$$

$$\delta \Psi_{R5} = \sum_n \delta \psi_{R5}^{(n)} f_{Rn} = \sum_n \left(\eta_R^{(n)} \partial_y \xi_{Rn} - b \sigma' \eta_R^{(n)} \xi_{Rn} \right). \quad (20)$$

Let us assume that the supersymmetry transformations for chiral modes of $\psi_5^{(n)}$ are

$$\delta \psi_{L5}^{(n)} = m_n \eta_L^{(n)}, \quad \delta \psi_{R5}^{(n)} = -m_n \eta_R^{(n)} \quad (21)$$

with m_n the four-dimensional mass of the n th gravitino KK mode, i.e., KK modes of Ψ_5 transform as the Goldstinos of the $\eta_{L,R}^{(n)}$ supersymmetry transformation. From Eqs. (19), (20), and (21), the relation for extra dimensional parts of Ψ_{L5} and Ψ_{R5} is

$$f_{Ln, Rn} = \frac{1}{m_n} (\pm \partial_y + b \sigma') \xi_{Ln, Rn}. \quad (22)$$

Therefore, the redefined gravitinos [60]

$$\tilde{\psi}_{L\mu}^{(n)} \equiv m_n \psi_{L\mu}^{(n)} - \partial_\mu \psi_{L5}^{(n)} + m_n \gamma_\mu \sum_k A_{kn} \frac{\psi_{R5}^{(k)}}{m_k}, \quad (23)$$

$$\tilde{\psi}_{R\mu}^{(n)} \equiv m_n \psi_{R\mu}^{(n)} + \partial_\mu \psi_{R5}^{(n)} - m_n \gamma_\mu \sum_k B_{kn} \frac{\psi_{L5}^{(k)}}{m_k} \quad (24)$$

are invariant under the supersymmetry transformations (17), (18), and (21), and they correspond to physical fields. While Ψ_5 is dependent on the gauge transformations and can be eliminated, which is the so-called super-Higgs mechanism.

Now we investigate the equations of motion for the redefined gravitino fields. The Rarita-Schwinger equation for the bulk gravitino field Ψ_M from the action (4) is

$$\Gamma^{[M} \Gamma^N \Gamma^{L]} D_N \Psi_L = a \sigma' \Gamma^{[M} \Gamma^{L]} \Psi_L, \quad (25)$$

and the boundary terms will vanish provided that

$$(\bar{\Psi}_{R\mu}\delta\Psi_L^\mu)|_{0,\bar{y}} = (\bar{\Psi}_{L\mu}\delta\Psi_R^\mu)|_{0,\bar{y}} = 0. \quad (26)$$

With the redefined fields (23) and (24), the equation (25) can be simplified to

$$\gamma^{[\lambda\mu\nu]}\partial_\mu\tilde{\psi}_{L\nu,R\nu}^{(n)} = m_n\gamma^{[\lambda\nu]}\tilde{\psi}_{R\nu,L\nu}^{(n)} \quad (27)$$

with the special parameters $b = \frac{1}{2}$ and $a = \frac{3}{2}$, which is the four-dimensional massive Rarita-Schwinger equation for the spin 3/2 field $\tilde{\psi}_\nu^{(n)}(x)$. And the KK modes $\xi_{Ln,Rn}(y)$ satisfy

$$\left[\pm\partial_y + \left(\frac{3}{2} \mp 1\right)\sigma'\right]\xi_{Ln,Rn} = m_n e^\sigma \xi_{Rn,Ln}, \quad (28)$$

which is consistent with the result in Ref. [60].

From above equations, we can obtain the solutions of the left- and right-handed zero modes:

$$\xi_{L0}(y) \propto e^{-\frac{1}{2}\sigma}, \quad \xi_{R0}(y) \propto e^{\frac{5}{2}\sigma}. \quad (29)$$

For the massive modes, we proceed with the coordinate and field transformations: $dy = e^{-\sigma}dz$, $\xi_{Ln,Rn}(y) = e^\sigma \bar{\xi}_{Ln,Rn}(z)$. Then the chiral coupling equations (28) can be rewritten as

$$\left(\pm\partial_z + \frac{3}{2}\partial_z\sigma\right)\bar{\xi}_{Ln,Rn} = m_n\bar{\xi}_{Rn,Ln}. \quad (30)$$

Thus, it is easy to see that KK modes $\bar{\xi}_{Ln,Rn}$ satisfy the following Schrödinger-like equations:

$$(-\partial_z^2 + V_{L,R}(z))\bar{\xi}_{Ln,Rn} = m_n^2\bar{\xi}_{Ln,Rn}, \quad (31)$$

where m_n is the mass of the gravitino KK mode $\tilde{\psi}_{L\nu,R\nu}^{(n)}$ and the effective potentials read

$$V_{L,R}(z) = \frac{9}{4}(\partial_z\sigma)^2 \mp \frac{3}{2}\partial_z^2\sigma. \quad (32)$$

The Schrödinger-like equations (31) can also be written as

$$Q^\dagger Q\bar{\xi}_{Ln} = m_n^2\bar{\xi}_{Ln}, \quad QQ^\dagger\bar{\xi}_{Rn} = m_n^2\bar{\xi}_{Rn} \quad (33)$$

with $Q = \partial_z + \frac{3}{2}\partial_z\sigma$, which indicates that there is no tachyon KK gravitino with negative mass square m_n^2 .

On the other hand, from Eqs. (22) and (28), we find that the conditions imposed on $f_{Ln,Rn}$ read

$$f_{Ln} = e^\sigma \xi_{Rn}, \quad f_{Rn} = e^\sigma \xi_{Ln} - \frac{2\sigma'}{m_n}\xi_{Rn}, \quad (34)$$

which is also consistent with the result given in Ref. [60]. Finally, we need the orthonormality conditions

$$\int dy e^{-\sigma} \xi_{Ln}(y) \xi_{Lk}(y) = \int dz \bar{\xi}_{Ln}(z) \bar{\xi}_{Lk}(z) = \delta_{nk}, \quad (35a)$$

$$\int dy e^{-\sigma} \xi_{Rn}(y) \xi_{Rk}(y) = \int dz \bar{\xi}_{Rn}(z) \bar{\xi}_{Rk}(z) = \delta_{nk}, \quad (35b)$$

$$\int dy e^{-\sigma} \xi_{Ln}(y) \xi_{Rk}(y) = \int dz \bar{\xi}_{Ln}(z) \bar{\xi}_{Rk}(z) = 0 \quad (35c)$$

to obtain the effective four-dimensional action for the massless and massive gravitino fields:

$$\begin{aligned} S_{\frac{3}{2}} &= \sum_n \int d^4x i \left(\bar{\tilde{\psi}}_{L\lambda}^{(n)} \gamma^{[\lambda\mu\nu]} \partial_\mu \tilde{\psi}_{L\nu}^{(n)} - m_n \bar{\tilde{\psi}}_{L\lambda}^{(n)} \gamma^{[\lambda\nu]} \tilde{\psi}_{R\nu}^{(n)} + \bar{\tilde{\psi}}_{R\lambda}^{(n)} \gamma^{[\lambda\mu\nu]} \partial_\mu \tilde{\psi}_{R\nu}^{(n)} - m_n \bar{\tilde{\psi}}_{R\lambda}^{(n)} \gamma^{[\lambda\nu]} \tilde{\psi}_{L\nu}^{(n)} \right) \\ &= \sum_n \int d^4x \left(\bar{\tilde{\psi}}_\lambda^{(n)} i \gamma^{[\lambda\mu\nu]} \partial_\mu \tilde{\psi}_\nu^{(n)} - m_n \bar{\tilde{\psi}}_\lambda^{(n)} i \gamma^{[\lambda\nu]} \tilde{\psi}_\nu^{(n)} \right). \end{aligned} \quad (36)$$

Note that the orthonormality conditions (35) are very essential to check whether the KK modes of the gravitino field can be localized on a brane.

Next we will consider the localization of the gravitino field in two concrete thin brane models (the RS1 and scalar-tensor models) and give its KK mass spectrum, which may be the signal of extra warped dimensions in future experiments.

III. LOCALIZATION OF THE GRAVITINO FIELD ON THE RS1 BRANE

In 1999, Randall and Sundrum presented the famous RS1 model [5] to solve the gauge hierarchy problem. In this section, we study the localization of the gravitino field on the RS1 brane with a compact extra dimension.

The line element describing two flat branes embedded in a five-dimensional spacetime is given by (1), where the fifth dimension is compactified on an orbifold S^1/Z_2 ($0 \leq y \leq \pi R$). For the RS1 brane model, the five-dimensional fundamental scale is the Planck scale $M_{\text{Pl}} \approx 10^{16} \text{ TeV}$ and the solution of the warped factor is

$$\sigma(y) = k|y| \quad (37)$$

with the anti-de Sitter curvature $k \sim M_{\text{Pl}}$, from which we have

$$\sigma'(y) = k\epsilon(y), \quad \sigma''(y) = 2k[\delta(y) - \delta(y - \pi R)]. \quad (38)$$




FIG. 1. The shapes of the warped factor $e^{-2\sigma}$ and chiral zero modes $\Psi_{L0} \equiv e^{-\frac{\sigma}{2}} \frac{\xi_{L0}}{\sqrt{k}}$ (the red dashed line), $\Psi_{R0} \equiv e^{-\frac{\sigma}{2}} \frac{\xi_{R0}}{\sqrt{k}}$ (the blue thick line).

With the solution (37), the left- and right-handed zero modes of the gravitino field (29) read

$$\xi_{L0} = \frac{1}{\sqrt{N_{L0}}} e^{-\frac{1}{2}k|y|}, \quad \xi_{R0} = \frac{1}{\sqrt{N_{R0}}} e^{\frac{1}{2}k|y|}, \quad (39)$$

where N_{L0} and N_{R0} are the normalization constants. As the result of the orthogonality conditions (35a) and (35b), the normalization constants $N_{L0} = \frac{1}{2k}(1 - e^{-2k\pi R}) \approx \frac{1}{2k}$, $N_{R0} = \frac{1}{4k}(e^{4k\pi R} - 1) \approx \frac{1}{4k}e^{4k\pi R}$, where we have considered that $kR \approx 12$ [5]. From Fig. 1, it seems that the left-handed zero mode is localized on the brane at $y = 0$ (UV brane), while the right-handed zero mode on the brane at $y = \pi R$ (IR brane). However, when considering the boundary conditions (26), we need that either ξ_{R0} is fixed on the boundaries with $\xi_{R0}|_{0,\pi R} = 0$ or ξ_{L0} is fixed. Thus we can obtain only one of the left- and right-handed zero modes on a brane. In fact this is how the four-dimensional chirality is recovered from five-dimensional bulk fermion fields, and it is the result of compactifying fermions on the S^1/Z_2 extra dimension. This property will be important to describe the Standard Model fermions since the left- and right-handed fermions transform differently under the electroweak gauge group [40].

For the massive KK modes, the effective potentials $V_{L,R}$ contain the second-order derivative of σ , which leads to the appearance of the delta functions. Thus, firstly, we can give the general solutions of the left- and right-handed KK gravitinos without their boundary behaviors, then consider their boundary conditions at $y = 0, \pi R$, respectively. Therefore, neglecting the term of σ'' and performing the transformations $z_n = \frac{m_n}{k}e^\sigma$, $\xi_{Ln,Rn}(y) = e^\sigma \bar{\xi}_{Ln,Rn}(z_n)$, the equation (28) will be transformed into the general ν -order Bessel equation:

$$z_n^2 \frac{d^2}{dz_n^2} \bar{\xi}_{Ln,Rn} + \left[z_n^2 - \frac{1}{k^2} \left(\frac{9}{4} \pm \frac{3}{2} \right) \sigma'^2 \right] \bar{\xi}_{Ln,Rn} = 0. \quad (40)$$

The general solution is $\bar{\xi}_{Ln,Rn} = z_n^\alpha Z_{\nu_\pm}(\lambda z_n^\beta)$, where the parameters are $\alpha = \frac{1}{2}$, $\beta = \lambda = 1$, $\nu_\pm = |\frac{3}{2} \pm \frac{1}{2}|$, and $Z_\nu(x)$ is the ν -order column function: $Z_\nu(x) = J_\nu(x) + b_\nu Y_\nu(x)$. The corresponding y -dependent left- and right-handed KK gravitinos read

$$\xi_{Ln}(y) = \frac{1}{N_n} e^{\frac{3}{2}\sigma} \left[J_2 \left(\frac{m_n}{k} e^\sigma \right) + b(m_n) Y_2 \left(\frac{m_n}{k} e^\sigma \right) \right], \quad (41)$$

$$\xi_{Rn}(y) = \frac{1}{N_n} e^{\frac{3}{2}\sigma} \left[J_1 \left(\frac{m_n}{k} e^\sigma \right) + b(m_n) Y_1 \left(\frac{m_n}{k} e^\sigma \right) \right], \quad (42)$$

where the coefficients N_n and $b(m_n)$ are arbitrary constants. These constants are the same for left- and right-handed KK modes since they are obeying Eq. (28), which is consistent with the result in [60].




FIG. 2. The shapes of three lowest chiral massive KK modes in the physical coordinate, $e^{-\frac{\sigma}{2}} \xi_{Ln,Rn}(y)$, and in the conformal coordinate, $\bar{\xi}_{Ln,Rn}(z)$, with the boundary conditions (43). The corresponding mass spectrum is given by $m_1 = 1.62508$ TeV (the dashed red lines), $m_2 = 2.97546$ TeV (the thickness blue lines), and $m_3 = 4.31474$ TeV (the thin green lines).

In the following, we focus on the boundary conditions imposing on $\xi_{Ln,Rn}$ at the fixed points $y = 0, \pi R$ and then give the mass spectrum of the four-dimensional KK gravitinos. It is clear that $\xi_{Ln,Rn}$ must be even (odd) as $\xi_{Rn,Ln}$ is odd (even) under the Z_2 symmetry, since the operator $[\pm \partial_y + (\frac{3}{2} \mp 1) \sigma']$ in Eq. (28) is odd. Firstly, considering that ξ_{Ln} is even, while ξ_{Rn} odd, we have

$$\left(\partial_y + \frac{\sigma'}{2} \right) \xi_{Ln} |_{0, \pi R} = 0, \quad \xi_{Rn} |_{0, \pi R} = 0. \quad (43)$$

From the above boundary conditions, $b(m_n) = -\frac{J_1(\frac{m_n}{k})}{Y_1(\frac{m_n}{k})}$, and the mass spectrum is determined by

$$b(m_n) = b(m_n e^{\pi k R}). \quad (44)$$

In the limit $m_n \ll k \sim M_{Pl}$ (the light KK modes compared with Planck mass) and $m_n \gg k e^{-k\pi R} \sim 1\text{TeV}$ (the KK modes with not too light mass compared with TeV scale), we obtain $b(m_n) \simeq \frac{m_n^2 \pi}{4k^2} \rightarrow 0$, and then

$$J_1\left(\frac{m_n}{k} e^{k\pi R}\right) \simeq \sqrt{\frac{2k e^{-k\pi R}}{\pi m_n}} \cos\left(\frac{m_n}{k} e^{k\pi R} - \frac{3\pi}{4}\right) \rightarrow 0. \quad (45)$$

Therefore the KK masses can be obtained approximately as

$$m_n \simeq \left(n + \frac{1}{4}\right) \pi k e^{-kR\pi} \sim 3 \left(n + \frac{1}{4}\right) \text{TeV} \quad (46)$$

with $n = 1, 2, 3, \dots$.




FIG. 3. The shapes of three lowest chiral massive KK modes in the physical coordinate, $e^{-\frac{\sigma}{2}} \xi_{Ln,Rn}(y)$, and in the conformal coordinate, $\bar{\xi}_{Ln,Rn}(z)$, with the boundary conditions (47). The corresponding mass spectrum is given by $m_1 = 2.17811$ TeV (the dashed red lines), $m_2 = 3.56988$ TeV (the thickness blue lines), and $m_3 = 4.92815$ TeV (the thin green lines).

Secondly, considering that ξ_{Rn} is even and ξ_{Ln} odd, we have the following boundary conditions

$$\left(-\partial_y + \frac{5\sigma'}{2}\right) \xi_{Rn} \big|_{0,\pi R} = 0, \quad \xi_{Ln} \big|_{0,\pi R} = 0. \quad (47)$$

So, the coefficient $b(m_n)$ is given as

$$b(m_n) = -\frac{J_2\left(\frac{m_n}{k}\right)}{Y_2\left(\frac{m_n}{k}\right)}, \quad (48)$$

which satisfies the expression $b(m_n) = b(m_n e^{\pi k R})$. In the limit $m_n \ll k$ and $m_n \gg k e^{-k\pi R} \sim 1\text{TeV}$, we have $b(m_n) \simeq \frac{\pi}{2} \left(\frac{m_n}{2k}\right)^4 \rightarrow 0$, and

$$J_2\left(\frac{m_n}{k} e^{k\pi R}\right) \simeq \sqrt{\frac{2k}{\pi m_n}} e^{-k\pi R} \cos\left(\frac{m_n}{k} e^{k\pi R} - \frac{5\pi}{4}\right) \rightarrow 0. \quad (49)$$

The KK masses are given approximately as

$$m_n \simeq \left(n + \frac{3}{4}\right) \pi k e^{-kR\pi} \sim 3 \left(n + \frac{3}{4}\right) \text{TeV}. \quad (50)$$

Three lowest chiral massive KK modes in the coordinates y and z are plotted in Fig. 2 (with the boundary conditions (43)) and Fig. 3 (with the boundary conditions (47)). From the point of view of the conformal coordinate z , the lower KK modes are not localized near any brane. However, as the statement in Ref. [64], from the point of view of the physical coordinate y , both the left- and right-handed lower KK modes are localized near the IR brane.

In this section, we have obtained the chiral zero modes and a set of massive KK modes of the gravitino field. There is only one of the localized chiral zero modes, either the left-handed zero mode near the UV brane or the right-handed one near the IR brane. Both the lower left- and right-handed massive KK modes are confined to the IR brane from

the point of view of the physical coordinate y . Here, we can see that even though two 3-branes are separated by a Planck scale distance $\pi R \sim 12l_{Pl}$, the mass gap of the KK modes is of about $\pi k e^{-kR\pi} \sim 3\text{TeV}$, which is very different from the KK theory [1] and ADD theory [3]. Furthermore, the left- and right-handed massive KK modes share the same mass spectrum but with different parity, and the lower ones are both localized near the IR brane.

IV. LOCALIZATION OF THE GRAVITINO FIELD ON THE THIN BRANES IN THE SCALAR-TENSOR THEORY

Another typical thin brane model is the one in the scalar-tensor gravity theory [43]. In this model, our world is confined to the positive tension brane rather than to the negative one in order to solve the gauge hierarchy problem, and the spacings of mass spectra for graviton and matter fields are very tiny (about 10^{-3}eV) [43, 46]. In the following, we investigate the KK modes of the gravitino field on the scalar-tensor brane.

The action of this brane system with a dilaton field nonminimally coupled to gravity reads [43]

$$S_5 = \frac{M_5^3}{2} \int dx^5 \sqrt{-g} e^{\lambda\phi} [\mathcal{R} - (3 + 4\lambda) \partial_M \phi \partial^M \phi], \quad (51)$$

where $\mathcal{R}$ is the five-dimensional scalar curvature, M_5 and ϕ are the five-dimensional scale of gravity and the dilaton field, respectively. Here, we also consider the static flat brane scenario with an S^1/Z_2 extra spatial dimension. The metric is assumed to be the one given in Eq. (1) with the physical extra dimension coordinate y , or

$$ds^2 = e^{-2\sigma} (\eta_{\mu\nu} dx^\mu dx^\nu + dz^2) \quad (52)$$

with the conformal extra dimension one $z \in [-z_b, z_b]$. Hence, the dilaton field ϕ can be supposed to be a function of the extra coordinate z , i.e., $\phi = \phi(z)$. There are two thin branes in this system, which are located at $z = 0$ (IR brane) and z_b (UV brane), contrary to the RS1 model. Furthermore, another difference from the RS1 model is that the five-dimensional fundamental scale is the electroweak scale, i.e., $M_5 = M_{EW} \sim 1\text{TeV}$. There are two kinds of brane solutions for different ranges of the coupling parameter λ , and they are physically equivalent from the view point of the four-dimensional graviton KK mass spectrum on the brane [43]. However, the KK mass spectra of matter fields are different for these two brane solutions [46].




FIG. 4. The shapes of the warped factor $e^{-2\sigma}$ and the right-handed zero mode $\Psi_{R0} \equiv \bar{\xi}_{R0}/\sqrt{\beta}$ in the conformal coordinate z for the first brane solution (54) in the scalar-tensor theory.

A. Solution 1

Firstly, we consider the solution satisfying the following equations

$$\dot{\phi} = -2\dot{\sigma}, \quad \ddot{\sigma} = (2\lambda + 3)\dot{\sigma}^2, \quad (\lambda \neq -3/2) \quad (53)$$

where the dot denotes the derivative with respect to the conformal coordinate z . The solution of the warped factor in this case reads

$$e^{-2\sigma} = (1 + \beta|z|)^{\frac{2}{3+2\lambda}}, \quad (54)$$




FIG. 5. The shapes of the left-handed zero mode $\Psi_{L0} \equiv \bar{\xi}_{L0}/\sqrt{\beta}$ in the conformal coordinate z for the first solution (54) in the scalar-tensor theory.

where the parameters $\beta > 0$, $\lambda < -3/2$. Its profile is plotted in Fig. 4(a), which is similar with the RS1 model. For convenience, we redefine the parameters $p_1 \equiv \frac{1}{3+2\lambda} (< 0)$ and $\tilde{z}_b \equiv 1 + \beta z_b (> 1)$. Then the physical coordinate y is given by

$$y = \int_0^z e^{-\sigma(w)} dw = \begin{cases} \frac{1}{\beta(1+p_1)} [(1 + \beta|z|)^{1+p_1} - 1] & \lambda \neq -2 \\ \frac{1}{\beta} \ln(1 + \beta|z|) & \lambda = -2 \end{cases}, \quad (55)$$

from which we can see that the bulk size is not dynamically fixed and there should be the Goldberger-Wise mechanism [69] to stabilize the size of the extra dimension. In particular, for $\lambda = -2$ the solution of the warp factor is the same as that in the RS1 model except that the energy scales k and β are different. Therefore, in the scalar-tensor brane model with $\lambda = -2$ the massless and massive modes of the gravitino field are the same with that obtained in the RS1 model formally, and the only difference is the mass gap.

The solutions of the left- and right-handed zero modes $\bar{\xi}_{L0,R0}$ for the brane solution (54) are

$$\bar{\xi}_{L0}(z) = \frac{1}{\sqrt{L_0}} (1 + \beta|z|)^{\frac{3}{2}p_1}, \quad (56)$$

$$\bar{\xi}_{R0}(z) = \frac{1}{\sqrt{R_0}} (1 + \beta|z|)^{-\frac{3}{2}p_1}. \quad (57)$$

From the orthogonality conditions (35), we have the normalized constants

$$L_0 = \begin{cases} \frac{1}{\beta} \ln \tilde{z}_b & \lambda = -3 \\ \frac{1}{(3p_1+1)\beta} (\tilde{z}_b^{3p_1+1} - 1) & \lambda \neq -3 \end{cases}, \quad R_0 = \begin{cases} \frac{1}{2\beta} \tilde{z}_b^2 & \lambda = -3 \\ \frac{1}{(1-3p_1)\beta} (\tilde{z}_b^{1-3p_1} - 1) & \lambda \neq -3 \end{cases}. \quad (58)$$

The profiles of $\frac{\bar{\xi}_{L0,R0}(z)}{\sqrt{\beta}}$ and $\frac{e^{-\frac{\sigma}{2}} \bar{\xi}_{L0,R0}(y)}{\sqrt{\beta}}$ with different values of λ ($\lambda < -\frac{3}{2}$) are shown in Figs. 4(b), 5, and 6, from which we can see that there are some interesting characters for the chiral zero modes:

- In the conformal coordinate:




FIG. 6. The shapes of the chiral zero modes $\Psi_{L0} \equiv e^{-\frac{\sigma}{2}} \xi_{L0} / \sqrt{\beta}$ (the blue lines) and $\Psi_{R0} \equiv e^{-\frac{\sigma}{2}} \xi_{R0} / \sqrt{\beta}$ (the red dashed lines) in the physical coordinate y for the first solution (54) in the scalar-tensor theory.

- The left-handed zero mode sharply decreases to zero with the increase of βz (see Fig. 5). This decreasing becomes slower and its value at $z = 0$ decreases with the increase of $|\lambda|$ but stops after $\lambda \leq -50$. So the left-handed zero mode is localized near the IR brane at $z = 0$ when the coupling parameter $-50 \lesssim \lambda < -\frac{3}{2}$, and it is distributed uniformly along the extra dimension when $\lambda \lesssim -1000$.
- The right-handed zero mode is almost zero near the location of the IR brane, and then sharply increases toward the UV brane when λ is closed to $-3/2$. This situation will happen for $-2.25 < \lambda < -\frac{3}{2}$, and the growth is linear as $\lambda = -2.25$ (see Fig. 4(b)), which means that this mode is localized near the UV brane when the coupling parameter satisfies $-2.25 \lesssim \lambda < -\frac{3}{2}$. With the continue decreasing of $\lambda < 50$, it will increase rapidly from the IR brane and rise slowly to the UV brane. For $\lambda \lesssim -1000$, the right-handed zero mode is distributed uniformly along the extra dimension, which is the same as the left-handed one.

• In the physical coordinate:

- The localization characters of the chiral zero modes are similar to the case in the conformal coordinate.

The result is that, under the boundary condition $\bar{\xi}_{Ln}(z)|_{0,z_b} = 0$, there is one chiral zero mode localized on the IR

brane only when $-50 \lesssim \lambda \leq -\frac{3}{2}$, which is similar to the case of the RS1 brane model. However, both the left- and right-hand zero modes become a constant when $\lambda \lesssim -1000$.




FIG. 7. The mass spectra of the gravitino field with the boundary condition $\bar{\xi}_{Ln}(z)|_{0,z_b} = 0$ for the first solution (54) in the scalar-tensor theory. The solid and dashed lines stand for the accurate (numerical) and approximate (analytical) mass spectra, respectively. The parameters are set to $\beta = 10^{12} \text{eV}$, $z_b = 10^4 \text{eV}^{-1}$, and $\lambda = -3.01, -3.5, -10, -100, -1000$.




FIG. 8. The shapes of three lowest chiral massive KK modes $\bar{\xi}_{Ln,Rn}$ with the boundary condition $\bar{\xi}_{Ln}(z)|_{0,z_b} = 0$ for the first solution (54) in the scalar-tensor theory. The parameter λ is set to -3.5 . The corresponding mass spectrum is $m_1 = 3.66232 \times 10^{-4} \text{eV}$ (the red dashed lines), $m_2 = 6.83487 \times 10^{-4} \text{eV}$ (the blue thickness lines), $m_3 = 9.98813 \times 10^{-4} \text{eV}$ (the black thin lines).

Next, we come to the massive modes of the gravitino field. The corresponding effective potentials $V_{L,R}^1$ (32) read as

$$V_{L,R}^{(1)} = \frac{3(3p_1 \mp 2)p_1\beta^2}{4(1 + \beta|z|)^2} \pm \frac{3\beta p_1[\delta(z) - \delta(z - z_b)]}{1 + \beta|z|}, \quad (59)$$

from which the general solutions of the left- and right-handed KK modes are given by

$$\bar{\xi}_{Ln}(z) = \frac{1}{N_n} \left(M_{0,-\lambda p_1}(\bar{z}_n) + c(m_n) W_{0,-\lambda p_1}(\bar{z}_n) \right), \quad (60)$$

$$\bar{\xi}_{Rn}(z) = \frac{1}{N_n} \left(M_{0,-p_2}(\bar{z}_n) + c(m_n) W_{0,-p_2}(\bar{z}_n) \right), \quad (61)$$

where $M_{a,b}(z)$ and $W_{a,b}(z)$ are two kinds of Whittaker functions, and

$$\bar{z}_n \equiv \frac{2im_n}{\beta}(1 + \beta|z|), \quad p_2 \equiv \frac{1 + 6p_1}{2}. \quad (62)$$

Note that two constants N_n and $c(m_n)$ are the same for left- and right-handed KK modes since $\bar{\xi}_{Ln,Rn}$ satisfy the first-order equations (30). The boundary conditions (26) will determine the constant $c(m_n)$ and mass spectrum m_n , while the orthonormal conditions (35) will fix N_n .




FIG. 9. The mass spectrum of the gravitino field with the boundary condition $\bar{\xi}_{Rn}(z)|_{0,z_b} = 0$ for the first solution (54) in the scalar-tensor theory. The parameters are set to $\beta = 10^{12}\text{eV}$, $z_b = 10^4\text{eV}^{-1}$, and $\lambda = -3.01, -3.5, -10, -100, -10000$.




FIG. 10. The shapes of three lowest chiral massive KK modes $\bar{\xi}_{Ln,Rn}$ with the boundary condition $\bar{\xi}_{Rn}(z)|_{0,z_b} = 0$ for the first solution (54) in the scalar-tensor theory. The parameter λ is set to -3.5 . The corresponding mass spectrum is $m_1 = 2.59517 \times 10^{-4}\text{eV}$ (the red dashed lines), $m_2 = 5.7144 \times 10^{-4}\text{eV}$ (the blue thickness lines), $m_3 = 8.84895 \times 10^{-4}\text{eV}$ (the black thin lines).

For the boundary conditions (26), we have two choices for the left- and right-handed massive KK modes: $\bar{\xi}_{Ln}(z)|_{0,z_b} = 0$, or $\bar{\xi}_{Rn}(z)|_{0,z_b} = 0$. We consider the first boundary condition $\bar{\xi}_{Ln}(z)|_{0,z_b} = 0$, i.e., the left-handed KK mode satisfies Dirichlet boundary condition, for which the boundary condition of right-handed KK mode should be $\partial_z \bar{\xi}_{Rn}(z)|_0 = -3\beta p_1 \bar{\xi}_{Rn}(0)$ and $\partial_z \bar{\xi}_{Rn}(z)|_{z_b} = \frac{3\beta p_1}{z_b} \bar{\xi}_{Rn}(z_b)$. The corresponding mass spectrum m_n is determined by

$$\frac{2^{-p_2} \Gamma(\lambda p_1) \sqrt{z_0} I_{\lambda p_1}(z_0)}{W_{0,\lambda p_1}(2z_0)} = \frac{1}{4\lambda p_1} \frac{M_{0,\lambda p_1}(\bar{z}_b)}{W_{0,\lambda p_1}(\bar{z}_b)}, \quad (63)$$

where $I_\nu(x)$ is the virtual variable ν -order Bessel function, $z_0 \equiv \frac{im_n}{\beta}$, and $\bar{z}_b \equiv \frac{2im_n}{\beta}(1 + \beta|z_b|)$. In the limit $\frac{m_n}{\beta} \ll 1$ and $1 \ll \beta z_b$, the left side of (63) is approximate to zero and the four-dimensional KK mass spectrum m_n is given by

$$m_n \simeq \frac{\beta x_n}{2(1 + \beta z_b)} \simeq \frac{x_n}{2z_b}, \quad (n = 1, 2, \dots) \quad (64)$$

where x_n satisfies $M_{0,\lambda p_1}(ix_n) = 0$. The accurate and approximate mass spectra of the gravitino field are shown in Fig. 7, which show that the approximate mass spectrum in this long wave limit is nearly close to the numerical result. The mass of the lightest massive KK mode decreases with the increase of $|\lambda|$. The mass gap of two adjoining KK modes ($\Delta m_n \simeq \frac{x_{n+1}-x_n}{2} 10^{-4} \text{eV}$) decreases with the level n and it will not change for heavy KK modes. And it also becomes smaller as decreasing the nonminimal coupling parameter λ .

For the second boundary condition $\bar{\xi}_{Ln}(z)|_{0,z_b} = 0$, it will result in $\partial_z \bar{\xi}_{Ln}(z)|_{0=0} = 3\beta p_1 \bar{\xi}_{Ln}(0)$, $\partial_z \bar{\xi}_{Ln}(z)|_{z_b=0} = -\frac{3\beta p_1}{\bar{z}_b} \bar{\xi}_{Ln}(z_b)$ and the mass spectrum m_n is determined by

$$\frac{2^{p_2-5\lambda p_1} \Gamma(1-\lambda p_1) \sqrt{z_0} I_{1-\lambda p_1}(z_0)}{W_{0,1-\lambda p_1}(2z_0)} = \frac{1}{4(1-\lambda p_1)} \frac{M_{0,1-\lambda p_1}(\bar{z}_b)}{W_{0,1-\lambda p_1}(\bar{z}_b)}. \quad (65)$$

The mass spectrum in this brane solution is shown in Fig. 9 for different values of λ . It can be seen that the mass of the lightest KK mode increases with $|\lambda|$, which is just opposite to the case of the first boundary condition $\bar{\xi}_{Ln}(z)|_{0,z_b} = 0$. The mass gap of two adjoin KK masses (Δm_n) changes with n and λ in the same way with the case of the boundary condition $\bar{\xi}_{Ln}(z)|_{0,z_b} = 0$. Here, we only plot the left- and right-handed massive gravitino KK modes with $\lambda = -3.5$ for both two kinds of boundary conditions in Figs. 8 and 10, respectively.

B. Solution 2




FIG. 11. The shapes of the warped factor $e^{-2\sigma}$ and the zero modes $\bar{\xi}_{L0}$ (the dashed lines), $\bar{\xi}_{R0}$ (the thick lines) for the second solution (67) in the scalar-tensor theory. The parameter λ is set to $\lambda = -0.98$ for the thin lines, $\lambda = -1$ for the thick lines.

Another solution of the warped factor in the scalar-tensor brane system satisfies the following equations

$$\dot{\sigma} = \frac{3+4\lambda}{6} \dot{\phi}, \quad \ddot{\sigma} = \left(\frac{3}{2} + \lambda\right) \dot{\phi}^2, \quad \left(\lambda \neq -\frac{3}{2} \text{ and } \lambda \neq -\frac{3}{4}\right) \quad (66)$$

and is given by [43]

$$e^{-2\sigma} = (1 + \beta|z|)^{\frac{2(3+4\lambda)}{9+6\lambda}}. \quad \left(-\frac{3}{2} < \lambda < -\frac{3}{4}\right) \quad (67)$$

The relation between the physical coordinate y and conformal one z becomes

$$y = \frac{9+6\lambda}{2(6+5\lambda)\beta} \left[(1 + \beta|z|)^{\frac{2(6+5\lambda)}{9+6\lambda}} - 1 \right], \quad (68)$$

and the chiral zero modes $\bar{\xi}_{L0,R0}$ read

$$\bar{\xi}_{L0}(z) = \frac{1}{\sqrt{L_0}} (1 + \beta|z|)^{\frac{3+4\lambda}{6+4\lambda}}, \quad (69)$$

$$\bar{\xi}_{R0}(z) = \frac{1}{\sqrt{R_0}} (1 + \beta|z|)^{-\frac{3+4\lambda}{6+4\lambda}}. \quad (70)$$

It is clear that the size of extra dimension is determined by the coupling parameter λ , which is similar to that in the first brane solution. The warp factor and chiral zero modes of the gravitino field are plotted in Fig. 11. Note that as



(a) With boundary condition $\bar{\xi}_{Rn}(z)|_{0,z_b} = 0$ (b) With boundary condition $\bar{\xi}_{Ln}(z)|_{0,z_b} = 0$

FIG. 12. The mass spectra of the gravitino field for two kinds of the boundary conditions in the scalar-tensor theory with the second solution (67). The parameters are set to $\beta = 10^{12} \text{eV}$, $z_b = 10^4 \text{eV}^{-1}$, and $\lambda = -1, -1.35, -1.42, -1.43, -1.45$.

the statements above, there will be only one chiral zero mode localized near one brane as considering the boundary conditions (26).

The effective potentials (32) with the brane solution (67) read

$$V_L^{(2)} = \frac{3(3p_1 - 2)p_1\beta^2}{4(1 + \beta|z|)^2} - (3p_1 - 2)\beta \frac{\delta(z) - \delta(z - z_b)}{1 + \beta|z|}, \quad (71a)$$

$$V_R^{(2)} = \frac{(3p_1 - 2)(3p_1 - 4)\beta^2}{4(1 + \beta|z|)^2} + (3p_1 - 2)\beta \frac{\delta(z) - \delta(z - z_b)}{1 + \beta|z|}. \quad (71b)$$

Note that the effective potentials (59) and (71) in the Shrödinger-like equations for two kinds of brane solutions are the same as $p_1 = 1/3$, which means that two brane solutions stand for the same geometric configuration of spacetime.

The general solutions of the left- and right-handed massive modes $\bar{\xi}_{Ln,Rn}$ are given by

$$\bar{\xi}_{Ln}(z) = \frac{1}{N_n} \left\{ M_{0,-\lambda p_1}(\bar{z}_n) + d(m_n) W_{0,-\lambda p_1}(\bar{z}_n) \right\}, \quad (72)$$

$$\bar{\xi}_{Rn}(z) = \frac{1}{N_n} \left\{ M_{0,-(1+\lambda p_1)}(\bar{z}_n) + d(m_n) W_{0,-(1+\lambda p_1)}(\bar{z}_n) \right\}, \quad (73)$$




FIG. 13. The shapes of three lowest chiral massive KK modes $\bar{\xi}_{Ln,Rn}$ with the boundary condition $\bar{\xi}_{Rn}(z)|_{0,z_b} = 0$ for the second solution (67) in the scalar-tensor theory. The parameter λ is set to -1 . The corresponding mass spectrum is $m_1 = 2.4048 \times 10^{-4} \text{eV}$ (the red dashed lines), $m_2 = 5.56445 \times 10^{-4} \text{eV}$ (the blue thickness lines), $m_3 = 8.69877 \times 10^{-4} \text{eV}$ (the black thin lines).




FIG. 14. The shapes of three lowest chiral massive KK modes $\bar{\xi}_{Ln,Rn}$ with the boundary condition $\bar{\xi}_{Ln}(z)|_{0,z_b} = 0$ for the second solution (67) in the scalar-tensor theory. The parameter λ is set to -1 . The corresponding mass spectrum is $m_1 = 3.83171 \times 10^{-4} \text{eV}$ (the red dashed lines), $m_2 = 7.01559 \times 10^{-4} \text{eV}$ (the blue thickness lines), $m_3 = 10.1735 \times 10^{-4} \text{eV}$ (the black thin lines).

where the constants N_n and $d(m_n)$ are the same for both the left- and right-handed KK modes. Performing the similar calculation with above subsection, the constant N_n will be fixed by the orthonormal conditions (35), and the mass spectrum m_n and $d(m_n)$ will be determined by the boundary conditions (26).

Firstly, we consider $\bar{\xi}_{Rn}|_{0,z_b} = 0$, i.e., the right-handed KK mode satisfies Dirichlet boundary condition, for which the left-handed KK mode at the boundaries satisfies $\partial_z \bar{\xi}_{Ln}(z)|_0 = -(2 - 3p_1)\beta \bar{\xi}_{Ln}(0)$ and $\partial_z \bar{\xi}_{Ln}(z)|_{z_b} = \frac{(2-3p_1)\beta}{z_b} \bar{\xi}_{Ln}(z_b)$. The corresponding mass spectrum m_n is determined by the following equation

$$\frac{M_{0,-(1+\lambda p_1)}(2z_0)}{W_{0,-(1+\lambda p_1)}(2z_0)} = \frac{M_{0,-(1+\lambda p_1)}(\bar{z}_b)}{W_{0,-(1+\lambda p_1)}(\bar{z}_b)}. \quad (74)$$

For the second boundary condition: $\bar{\xi}_{Ln}|_{0,z_b} = 0$, the right-handed KK modes at the boundaries satisfy $\partial_z \bar{\xi}_{Rn}(z)|_0 = (2 - 3p_1)\beta \bar{\xi}_{Rn}(0)$, $\partial_z \bar{\xi}_{Rn}(z)|_{z_b} = -\frac{(2-3p_1)\beta}{z_b} \bar{\xi}_{Rn}(z_b)$, and the mass spectrum m_n is determined by

$$\frac{M_{0,-\lambda p_1}(2z_0)}{W_{0,-\lambda p_1}(2z_0)} = \frac{M_{0,-\lambda p_1}(\bar{z}_b)}{W_{0,-\lambda p_1}(\bar{z}_b)}. \quad (75)$$

The numerical result of the gravitino mass spectra for two kinds of boundary conditions is listed in Fig. 12, which shows that the mass m_1 of the lightest massive KK mode increases with the coupling parameter $|\lambda|$ for both two boundary conditions. Furthermore, the mass gap Δm_n of two adjacent KK massive modes becomes smaller with the decrease of $|\lambda|$ and it will not change for the heavy chiral KK modes. These results are not all the same with that in the first brane solution. From Figs. 7, 9, and 12, we know that these two brane solutions are not physically equivalent from the view of the gravitino mass spectrum. Here, we only show three lowest chiral massive KK modes with $\lambda = -1$ for different boundary conditions in Figs. 13 and 14, respectively.

V. CONCLUSION

In this paper, we have investigated the massless and massive KK modes of a bulk massive gravitino field in the RS1 and scalar-tensor thin brane models. The localization and mass spectrum the gravitino field on a brane are obtained for both models.

In the case of the RS1 model, the left- and right-handed zero modes (four-dimensional massless gravitinos) are localized near the Planck brane (UV brane at $y = 0$) and TeV brane (IR brane at $y = y_b$), respectively. However, there is only one massless chiral gravitino survived as the result of the boundary conditions, which is consistent with Refs. [35, 61]. The left- and right-handed massive KK modes share the same mass spectrum and the mass gap is of about 3TeV. The localization behaviors of the left- and right-handed lower massive KK modes in the conformal and physical coordinates are different. From the point of view of the conformal coordinate, the lower massive KK modes are not localized near any brane. However, they are confined near the IR brane from the point of view of the physical one. These behaviors of a gravitino field are similar to the Dirac fermion field in the RS1 model.

For the scalar-tensor thin brane model, there are two brane solutions. The left- and right-handed zero modes are localized near the IR brane (at $y = 0$) and UV brane (at $y = y_b$), respectively, if the coupling parameter $|\lambda|$ is not too large for the first brane solution and not close to 3/4 for the second one. The mass spectra and localization properties of the gravitino field for two brane solutions are different. The mass gap is very small and is of about 10^{-4}eV . For the first brane solution, the mass m_1 of the lightest massive gravitino decreases with the coupling $|\lambda|$ for the boundary conditions $\bar{\xi}_{Ln}(z)|_{0,z_b} = 0$, while it will increases with $|\lambda|$ for $\bar{\xi}_{Rn}(z)|_{0,z_b} = 0$. For the second one, the mass m_1 increases with the coupling $|\lambda|$ for both boundaries. The lower massive KK modes are not localized near any brane except for the first brane solution with $\bar{\xi}_{Rn}(z)|_{0,z_b} = 0$, for which the left-handed KK modes are confined near the IR brane. These results show that the two brane solutions are not physically equivalent.

In the future work, we should introduce interaction between gravitino and other fields and consider the localization and resonances of the gravitino field on thick branes.

ACKNOWLEDGMENTS

We would like to thank Dr. Yuan Zhong for helpful discussion. This work was supported by the National Natural Science Foundation of China (Grant No. 10905027, No. 11375075, and No. 11522541), and the Fundamental Research Funds for the Central Universities (Grant No. lzujbky-2015-jl1). Y. Zhong was supported by the scholarship granted by the Chinese Scholarship Council (CSC).

-
- [1] T. Kaluza, Sitzungsber. Preuss. Akad. Wiss. Berl. Math. Phys. K. **1**, 966 (1921).
 - [2] O. Klein, Z. Phys. **37**, 895 (1926).
 - [3] N. Arkani-Hamed, S. Dimopoulos, and G. R. Dvali, Phys. Lett. B **429**, 263 (1998).
 - [4] I. Antoniadis, N. Arkani-Hamed, S. Dimopoulos, and G. R. Dvali, Phys. Lett. B **436**, 257 (1998).
 - [5] L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, 3370 (1999).
 - [6] L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, 4690 (1999).
 - [7] I. Antoniadis, K. Benakli, and M. Quiros, Phys. Lett. B **460**, 176 (1999).
 - [8] DØ collaboration, B. Abbott *et al.*, Phys. Rev. Lett. **86**, 1156 (2001).
 - [9] S. Nussinov and R. Shrock, Phys. Rev. Lett. **88**, 171601 (2002).
 - [10] A. A. Pankov and N. Paver, Phys. Rev. D **72**, 035012 (2005).
 - [11] CDF collaboration, A. Abulencia *et al.*, Phys. Rev. Lett. **97**, 171802 (2006).
 - [12] DELPHI collaboration, J. Abdallah *et al.*, Eur. Phys. J. C **60**, 17 (2009).
 - [13] R. Franceschini, P. P. Giardino, G. F. Giudice, and A. Strumia, J. High Energy Phys. **1105** (2011) 092.
 - [14] V. S. Baste-Gonzalez, A. Esmaili, and O. L. G. Peres, Phys. Lett. B **718**, 1020 (2013).
 - [15] H. Sun, Eur. Phys. J. C **74**, 2977 (2014).
 - [16] K. Cheung, Phys. Rev. Lett. **88**, 221602 (2002).
 - [17] A. Casanova and E. Spallucci, Classical Quantum Gravity **23**, R45 (2006).
 - [18] T. G. Rizzo, Phys. Lett. B **647**, 43 (2007).
 - [19] S. C. Park, H. S. Song, and J. H. Song, Phys. Rev. D **65**, 075008 (2002).
 - [20] X.-Z. Li, W.-G. Ma, R. Y. Zhang, and L. Guo, Phys. Rev. D **87**, 056008 (2013).
 - [21] I. Sahin, M. Koksul, S. Inan, and A. Billur, Phys. Rev. D **91**, 035017 (2015).
 - [22] C. D. Hoyle, U. Schmidt, B. R. Heckel, E. G. Adelberger, J. H. Gundlach, D. J. Kapner, and H. E. Swanson, Phys. Rev. Lett. **86**, 1418 (2001).

- [23] C. D. Hoyle, D. J. Kapner, B. R. Heckel, E. G. Adelberger, J. H. Gundlach, U. Schmidt, and H. E. Swanson, Phys. Rev. D **70**, 042004 (2004).
- [24] E. G. Adelberger, B. R. Heckel, S. A. Hoedl, C. D. Hoyle, D. J. Kapner, and A. Upadhye, Phys. Rev. Lett. **98**, 131104 (2007).
- [25] D. J. Kapner, T. S. Cook, E. G. Adelberger, J. H. Gundlach, B. R. Heckel, C. D. Hoyle, and H. E. Swanson, Phys. Rev. Lett. **98**, 021101 (2007).
- [26] C.-G. Shao, Y.-J. Tan, W.-H. Tan, S.-Q. Yang, J. Luo, and M. E. Tobar, Phys. Rev. D **91**, 102007 (2015).
- [27] C. Csaki, J. Erlich, T. J. Hollowood, and Y. Shirman, Nucl. Phys. B **581**, 309 (2000).
- [28] M. Gremm, Phys. Lett. B **478**, 434 (2000).
- [29] B. Bajc and G. Gabadadze, Phys. Lett. B **474**, 282 (2000).
- [30] T. Shiromizu, K.-I. Maeda, and M. Sasaki, Phys. Rev. D **62**, 024012 (2000).
- [31] I. Oda, Phys. Lett. B **508**, 96 (2001).
- [32] B. Mukhopadhyaya, S. Sen, and S. SenGupta, Phys. Rev. D **70**, 066009 (2004).
- [33] A. Chatterjee and P. Majumdar, Phys. Rev. D **72**, 066013 (2005).
- [34] A. Melfo, N. Pantoja, and J. D. Tempo, Phys. Rev. D **73**, 044033 (2006).
- [35] Y.-X. Liu, L. Zhao, and Y.-S. Duan, J. High Energy Phys. **0704** (2007) 097.
- [36] A. de Souza Dutra, J. de Faria, A. C. Amaro, and M. Hott, Phys. Rev. D **78**, 043526 (2008).
- [37] M. Tahim, W. Cruz, and C. Almeida, Phys. Rev. D **79**, 085022 (2009).
- [38] C. A. S. Almeida, M. M. Ferreira, Jr. A. R. Gomes, and R. Casana, Phys. Rev. D **79**, 125022 (2009).
- [39] D. Bazeia, A. Gomes, and L. Losano, Int. J. Mod. Phys. A **24**, 1135 (2009).
- [40] T. Gherghetta, *TASI Lectures on a Holographic View of Beyond the Standard Model Physics*, arXiv:1008.2570.
- [41] R. Guerrero, A. Melfo, N. Pantoja, and R. O. Rodriguez, Phys. Rev. D **81**, 086004 (2010).
- [42] R. Landim, G. Alencar, M. Tahim, and R. Costa Filho, J. High Energy Phys. **1108** (2011) 071.
- [43] K. Yang, Y.-X. Liu, Y. Zhong, X.-L. Du, and S.-W. Wei, Phys. Rev. D **86**, 127502 (2012).
- [44] C.-E. Fu, Y.-X. Liu, K. Yang, and S.-W. Wei, J. High Energy Phys. **1210** (2012) 060.
- [45] W. Cruz, R. Maluf, and C. Almeida, Eur. Phys. J. C **73**, 2523 (2013).
- [46] Q.-Y. Xie, Z.-H. Zhao, Y. Zhong, J. Yang, and X.-N. Zhou, J. Cosmol. Astropart. Phys. **1503** (2015) 014.
- [47] Z.-H. Zhao, Q.-Y. Xie, and Y. Zhong, Classical Quantum Gravity **32**, 035020 (2015).
- [48] E. J. Chun, H. B. Kim, and J. E. Kim, Phys. Rev. Lett. **72**, 1956 (1994).
- [49] M. Viel, J. Lesgourgues, M. G. Haehnelt, S. Matarrese, and A. Riotto, Phys. Rev. D **71**, 063534 (2005).
- [50] F. D. Steffen, J. Cosmol. Astropart. Phys. **0609** (2006) 001.
- [51] G. Panotopoulos, J. Cosmol. Astropart. Phys. **0705** (2007) 016.
- [52] R. Ding and Y. Liao, J. High Energy Phys. **1204** (2012) 054.
- [53] K. G. Savvidy and J. D. Vergados, Phys. Rev. D **87**, 075013 (2013).
- [54] J. L. Feng, M. Kamionkowski, and S. K. Lee, Phys. Rev. D **82**, 015012 (2010).
- [55] S. Shirai and T. T. Yanagida, Phys. Lett. B **680**, 351 (2009).
- [56] P. Arnold, P. Szepletowski, and D. Vaman, Phys. Rev. D **89**, 046001 (2014).
- [57] O. P. F. Piedra, Int. J. Mod. Phys. D **20**, 93 (2011).
- [58] A. Yale and R. B. Mann, Phys. Lett. B **673**, 168 (2009).
- [59] M. Yu. Khlopov, A. Barrau, and J. Grain, Classical Quantum Gravity **23**, 1875 (2006).
- [60] T. Gherghetta and A. Pomarol, Nucl. Phys. B **602**, 3 (2001).
- [61] I. Oda, Prog. Theor. Phys. **105**, 667 (2001).
- [62] H. M. Lee and A. Papazoglou, Nucl. Phys. B **792**, 166 (2008).
- [63] J. L. Hewett and D. Sadri, Phys. Rev. D **69**, 015001 (2004).
- [64] T. Gherghetta and A. Pomarol, Nucl. Phys. B **586**, 141 (2000).
- [65] E. Bergshoeff, R. Kallosh, and A. Van Proeyen, J. High Energy Phys. **0010** (2000) 033.
- [66] R. Altendorfer, J. Bagger, and D. Nemeschansky, Phys. Rev. D **63**, 125025 (2001).
- [67] N. Alonso-Alberca, P. Meessen, and T. Ortin, Phys. Lett. B **482**, 400 (2000).
- [68] A. Falkowski, Z. Lalak, and S. Pokorski, Phys. Lett. B **491**, 172 (2000).
- [69] W. D. Goldberger and M. B. Wise, Phys. Rev. Lett. **83**, 4922 (1999).